

Healthy Nutrition and Fermented Animal-Sourced Foods as a Complementary One Health Strategy: Effects on Foodborne Pathogens and Intestinal Host Defense

 Serhat Al¹

¹Department of Veterinary Public Health, Erciyes University, Veterinary Faculty, Kayseri, Türkiye

ABSTRACT

Foodborne diseases remain a major global public health challenge at the interface of human, animal, and environmental health, emphasizing the need for integrated preventive strategies within the One Health framework. Although animal-sourced foods (ASFs) are essential components of human nutrition, they also serve as important reservoirs and transmission vehicles for foodborne pathogens. Conventional food safety interventions primarily target microbial contamination within food systems but often fail to adequately address host susceptibility, intestinal resilience, and microbiota-mediated defense mechanisms. This review highlights fermented ASFs, nutraceuticals, and nonviable microbial products as complementary One Health interventions that can simultaneously influence pathogen ecology and host defense. Fermentation-derived compounds, including organic acids, bacteriocins, bioactive peptides, and microbial metabolites, exert antimicrobial effects against foodborne pathogens while modulating epithelial barrier integrity, inflammatory signaling pathways, gut microbiota composition, and colonization resistance. Particular emphasis is placed on the mechanistic roles of these functional components in regulating intestinal immunity, including Toll-like receptor-associated signaling, NF- κ B-mediated responses, and epithelial protection. The review also discusses multidisciplinary surveillance approaches, the translational relevance of dietary strategies for reducing reliance on antimicrobials, and current challenges related to standardization, safety, and clinical validation. Collectively, fermented ASFs and functional nutritional approaches should not be considered passive dietary components but biologically active tools with the potential to enhance food safety, strengthen gastrointestinal host defense, and support sustainable disease prevention across the interconnected One Health framework.

Keywords: Animal-sourced foods, fermentation, functional compounds, host defense, one health.



Cite this article as:

Al S. Healthy Nutrition and Fermented Animal-Sourced Foods as a Complementary One Health Strategy: Effects on Foodborne Pathogens and Intestinal Host Defense. J Clin Pract Res 2026;48(0):000–00.

Address for correspondence:

Serhat Al.
Department of Veterinary Public Health, Erciyes University, Veterinary Faculty, Kayseri, Türkiye
Phone: +90 543 325 87 41
E-mail: serhatal@erciyes.edu.tr

Submitted: 08.05.2026

Accepted: 11.06.2026

Available Online: 18.06.2026

Erciyes University Faculty of Medicine Publications - Available online at www.jcpres.com

INTRODUCTION

Foodborne diseases remain a persistent global challenge at the intersection of human, animal, and environmental health, closely aligning with the core principles of the One Health approach. Animal-sourced foods (ASFs) are essential components of human nutrition.¹ However, ASFs can serve as both



Copyright © Author(s)
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

major reservoirs and transmission vehicles for key foodborne pathogens, such as *Salmonella enterica*, *Listeria monocytogenes*, and *enterohemorrhagic Escherichia coli*.² These products are inherently nutrient-rich environments that can support microbial survival and growth if not properly handled. Although conventional control strategies, such as hygiene interventions, antimicrobial use, and advanced processing technologies, have improved safety outcomes, pathogens continue to occur in animal production systems and postharvest food chains. The increasing prevalence of antimicrobial resistance and the complexity of farm-to-fork transmission dynamics emphasize the need for innovative strategies that simultaneously target both the food matrix and the host.

Although ASFs pose risks due to the presence of zoonotic microorganisms, they also contain many functional components that provide benefits beyond their nutritional value. Animal-sourced fermented foods represent a compelling intersection of nutrition, microbiology, and food safety. Traditional products, such as fermented dairy foods (e.g., yogurt, kefir, and cheese) and fermented meats (e.g., dry-cured and ripened products), are biological systems shaped by complex microbial communities. These matrices generate a wide spectrum of bioactive metabolites during fermentation, transforming raw animal substrates into functional foods with enhanced safety and health-promoting properties.³ The starter microorganisms responsible for food fermentation also exhibit probiotic activities that can suppress pathogenic bacteria in food while simultaneously benefiting the host within the gastrointestinal system.⁴

Beyond their role in food preservation, fermented ASFs actively contribute to intestinal host defense. In addition to competing for resources within food matrices, metabolites produced during fermentation, including organic acids, bacteriocins, short-chain fatty acids, and bioactive peptides, can directly inhibit pathogens by acidifying the environment and destabilizing bacterial membranes.⁵ Upon consumption, these compounds interact with the intestinal epithelium, influencing barrier integrity and modulating immune signaling pathways, including pathogen recognition receptors and cytokine and chemokine release.⁶ This dual functionality, acting in both food matrices and the host response, positions fermented ASFs as uniquely effective tools for mitigating foodborne infections.

In this review, healthy nutrition and fermented ASFs are presented as complementary One Health strategies that bridge food safety and host health. The review focuses on their capacity to control foodborne pathogens at the source while enhancing intestinal defense mechanisms through modulation of the microbiota and immune regulation. By integrating insights from food microbiology, animal

production systems, and host–pathogen interaction studies, this review aims to redefine healthy nutrition and animal-sourced fermented foods not as passive dietary components but as active, biologically driven interventions with the potential to reduce disease burden across interconnected human, animal, and environmental systems.

BURDEN OF FOODBORNE PATHOGENS

Foodborne diseases remain a major global cause of outbreaks and contribute to substantial morbidity, mortality, and economic loss. Bacterial pathogens remain among the primary etiological agents of severe foodborne diseases, including *Salmonella enterica*, *Listeria monocytogenes*, and *Escherichia coli* O157:H7. Viral and parasitic agents, such as *norovirus* and *Toxoplasma gondii*, also require close monitoring to maintain food safety. These organisms are responsible for a wide spectrum of disease outcomes, ranging from self-limiting gastroenteritis to systemic infections, invasive disease, and life-threatening complications, particularly in vulnerable populations, including infants, older adults, and immunocompromised individuals. Despite advances in surveillance and food safety management systems, the incidence of foodborne illness remains high, reflecting the complexity of pathogen transmission across the entire food chain.⁷

Economic Consequences of Foodborne Illness

According to the World Health Organization (WHO), foodborne diseases remain a major global health concern, causing approximately 600 million cases and 420,000 deaths annually, with nearly 30% of these deaths occurring in children younger than 5 years. The burden extends beyond mortality, accounting for an estimated 33 million disability-adjusted life years (DALYs) lost each year.⁸ This substantial burden also translates into significant economic losses due to healthcare costs, reduced productivity, and disruptions across the food supply chain. Moreover, the disproportionate impact on vulnerable populations and low-resource settings highlights the urgent need for integrated preventive strategies aligned with the One Health framework. The United States Department of Agriculture (USDA) Economic Research Service (ERS) provides comprehensive data on the economic burden of foodborne illnesses in the United States, estimating the total cost at \$74.7 billion in 2023. These estimates incorporate medical treatment costs, lost wages, and short- and long-term health outcomes. Although serious illnesses caused by specified pathogens account for only 20% of total cases, they account for 60% of the total financial impact, with nontyphoidal *Salmonella* having the highest total cost in the United States at \$17 billion.⁹

Social Consequences of Foodborne Illness

The social burden of foodborne illness, in addition to its economic burden, must be re-evaluated in terms of human

welfare. The social costs of foodborne pathogens in vulnerable populations are substantial; beyond mortality, they directly affect education, social life, and acute clinical conditions, as well as complications such as Guillain-Barré syndrome (GBS) and hemolytic uremic syndrome (HUS), which can have long-term effects on quality of life. In addition to these direct effects, the psychological impact of foodborne pathogens on perceptions of healthy nutrition is undeniable. Food safety incidents affect public trust in food systems and regulatory authorities, altering consumer behavior and dietary patterns.¹⁰ Growing concerns about food safety and quality are also shaping food preferences. Evidence suggests that foodborne outbreaks can reduce the consumption of certain foods, although these effects vary by context and food type.¹¹ From this perspective, foodborne illnesses require not only symptomatic treatment but also the development of pathogen surveillance systems, integrated complementary approaches to healthy nutrition in the management of gastrointestinal diseases, and the use of preventive strategies during healthy phases of life.

ONE HEALTH INTEGRATION AND MULTIDISCIPLINARY SURVEILLANCE

Strategies aligned with the One Health paradigm emphasize the interrelationships among food, animal, and human health in addressing the persistent burden of foodborne diseases. The lack of One Health coordination impedes the effective integration of environmental, animal, and human health data, resulting in fragmented surveillance and slower responses to zoonotic threats. Disciplines often operate in isolation, with poor interdisciplinary communication, inconsistent data-sharing protocols, and a lack of standardized formats, leading to duplicated efforts and incomplete outbreak insights. Additional barriers include limited funding, weak governance, and insufficient multidisciplinary training. Real-time surveillance and response within a One Health framework require multidisciplinary expertise spanning epidemiology, veterinary medicine, genomics, bioinformatics, and laboratory practice. The value of this approach lies not only in integrating data across disciplines but also in enabling joint analyses that strengthen early detection and coordinated responses. By supporting operational data integration, real-time pathogen surveillance, and the clinical management of foodborne diseases, this framework facilitates collaborative disease prevention and control across human, animal, and environmental health systems.¹²

Interdisciplinary Collaboration

Effective control of foodborne outbreaks and disease management, as well as the successful implementation of One Health strategies, requires close and sustained collaboration across disciplines. The interface between

animal and human health is particularly evident in the epidemiology of zoonotic pathogens that originate in animal reservoirs and are transmitted to humans through the food chain. Veterinary specialists play a critical role in monitoring animal health, implementing biosecurity measures, and controlling pathogen prevalence at the pre- and postharvest levels,¹³ while medical professionals are responsible for the clinical diagnosis, treatment, and management of infections and outbreaks.¹⁴ Food safety also requires the coordinated involvement of food microbiologists, food safety specialists, public health authorities, and regulatory agencies. Without coordinated efforts among these disciplines, interventions remain fragmented and less effective.

Benefits of Interdisciplinary Collaboration

Collaboration enables the integration of surveillance systems, allowing early detection and rapid response to emerging foodborne threats. Data sharing between veterinary and medical professionals facilitates source attribution, outbreak tracing, and risk assessment, thereby improving the precision of intervention strategies. Moreover, joint research efforts can advance the understanding of host–pathogen interactions across species, particularly regarding immune responses, microbiota dynamics, and antimicrobial resistance.

Interdisciplinary collaboration in food safety supports the design of prevention strategies that extend from the farm to the clinic. Veterinary expertise in pathogen control supports approaches to reduce zoonotic occurrence before food products reach consumers. Simultaneously, medical insights into human immunity and disease susceptibility can inform dietary recommendations and functional nutrition strategies to enhance host defense. This knowledge exchange is essential for aligning food safety interventions with public health outcomes. Ultimately, integrating veterinary and medical sciences strengthens the One Health approach by bridging gaps among animal production, food safety, and human health.¹⁵ Coordinated efforts not only improve the effectiveness of pathogen control strategies but also help reduce antimicrobial resistance, enhance surveillance capacity, and promote sustainable health systems. This collaborative framework is indispensable for addressing the complex, interconnected challenges that foodborne diseases pose to the public. Although traceable pathogen control from farm to fork is well developed with modern management systems, food safety remains a critical public health challenge in both developed and developing countries. Therefore, strengthening nonspecific host defense barriers in the gastrointestinal system through healthy nutrition requires collaboration among related scientific fields, such as human medicine, nutrition, and dietetics, all of which are relevant to alimentary health.




Figure 1. Interactions among fermented animal-sourced foods, gut microbiota, foodborne pathogens, and the host intestinal environment.

ASF: Animal-sourced food; SCFA: Short-chain fatty acid; STEC: Shigatoxigenic *Escherichia coli*. (Gastrointestinal illustration was supplied from BioArt (<https://bioart.niaid.nih.gov/>)).

ROLE OF DIETARY COMPOUNDS IN MODULATING GASTROINTESTINAL DISEASE

Diet is a primary ecological determinant of host–microbe interactions and exerts system-wide effects that directly influence disease susceptibility across human, animal, and environmental niches. Beyond its fundamental role in energy provision, diet regulates the composition and metabolic activity of the gut microbiota, thereby enhancing colonization resistance against enteric pathogens. Nutrient availability and substrate diversity selectively shape microbial communities, influencing the production of metabolites such as short-chain fatty acids (SCFAs), secondary bile acids, and antimicrobial peptides. These metabolites act as key signaling molecules that reinforce epithelial barrier integrity, regulate mucosal immunity, and suppress pathogen growth.¹⁶ Consequently, dietary patterns rich in bioactive and fermentable substrates can shift the intestinal ecosystem toward a protective state, whereas nutrient imbalances or highly processed diets may predispose hosts to dysbiosis and increased infection risk. Integrating functional nutrition, fermented foods, nutraceuticals, and nonviable microbial products into human

dietary practices represents a rational pathway for reducing the burden of infectious diseases while promoting sustainable health across interconnected biological systems. Interactions among fermented ASFs, gut microbiota, foodborne pathogens, and the host intestinal environment are shown in Figure 1.

Fermented ASFs and Functional Compounds

Produced through controlled microbial activity, fermented ASFs, such as dairy and meat products, are not only preserved and stabilized but also enriched with bioactive compounds that enhance their functional value. During fermentation, lactic acid bacteria and other microbial consortia metabolize proteins, lipids, and carbohydrates to generate a wide spectrum of functional molecules, including organic acids, bacteriocins, short-chain fatty acids, vitamins, and bioactive peptides.¹⁷ Fermented foods help reduce the burden of foodborne pathogens through both direct and indirect mechanisms. During fermentation, the production of organic acids, bacteriocins, and other antimicrobial compounds can suppress or inhibit pathogens. These constraints are particularly relevant in ASFs and have led to the development of technological approaches

Table 1. Bioactive components and mechanistic health effects of fermented ASFs

ASF	Major functional components	Key mechanistic actions	Ref
Yogurt	ACE-inhibitory peptides	Antihypertensive activity mediated by bioactive peptide-induced inhibition of ACE	19
Acidophilus-fortified milk	Organic acids, bacteriocin-like compounds	Recognition through TLR signaling, inhibition of allergic responses, and regulation of inflammatory pathways, such as NF- κ B signaling	20
Milk	Casein-derived peptides	Enhanced MUC5AC gene expression and mucin secretion	21
Milk	Sphingomyelin	Modulation of lipid metabolism and gut microbial composition	22
Fermented milk	Fermentation metabolites	Modulation of gut microbiota composition and regulation of GALT	23
Kefir	EPS	Modulation of the microbiota	24
Koumiss	ACE-inhibitory peptides	Antihypertensive activity mediated by bioactive peptide-induced inhibition of ACE	25
Fermented skim milk	Functional peptide fractions	Antioxidant, antihypertensive, and immunomodulatory activities	26
Dry-cured ham	ACE-inhibitory peptides	Antihypertensive activity mediated by bioactive peptide-induced inhibition of ACE	27
Fermented meat	GABA	Neurotransmitter produced during fermentation with blood pressure-lowering effects	28
Probiotic salami	SCFAs	Regulation of IEC function through mechanisms including histone deacetylase inhibition	29
Dry-fermented sausage	ACE-inhibitory peptides	Antihypertensive activity mediated by bioactive peptide-induced inhibition of ACE	30
Salami	CLA	Anticarcinogenic activity	31

ASF: Animal-sourced food; CLA: Conjugated linoleic acid; SCFA: Short-chain fatty acid; GABA: γ -aminobutyric acid; EPS: Exopolysaccharide; ACE: Angiotensin-converting enzyme; TLR: Toll-like receptor; IEC: Intestinal epithelial cell; GALT: Gut-associated lymphoid tissue; Ref: Reference.

to ensure their safety, as conventional decontamination methods may have limited penetration into complex tissues, such as lymph nodes in meat, and may not adequately address internalized pathogens. One study examined the effectiveness of carcass vascular rinsing with organic acids and acid-based antimicrobial interventions in reducing *Salmonella* Enteritidis in the lymph nodes of experimentally infected meat animals.¹⁸ The implementation of organic acids, which can be end products of beneficial microorganisms, such as lactic acid bacteria with probiotic and starter culture properties, into new technologies is important for food safety.

The consumption of fermented foods supports intestinal health by promoting beneficial microbiota, strengthening epithelial barrier function, and modulating immune responses, thereby reducing host susceptibility to enteric infections. Their role is particularly relevant in the context of increasing antimicrobial resistance and the limitations of conventional food safety strategies, as they offer natural and sustainable benefits that enhance both microbial control and host resilience. Furthermore, fermentation processes can

contribute to sustainability by improving food preservation, reducing waste, and enabling the more efficient use of food resources. Some key bioactive components of fermented ASFs and their mechanistic effects on host physiology and pathogen control are summarized in Table 1.^{19–31}

Fermented foods are integral components of traditional dietary systems across nearly all global cultures, reflecting long-standing practices of food preservation, sensory enhancement, and microbial diversity. These culturally embedded foods contribute not only to nutritional intake but also to social eating behaviors, communal food sharing, and dietary continuity, which are increasingly recognized as determinants of long-term health outcomes. The sustained consumption of traditional fermented foods may support healthier dietary patterns by reinforcing culturally accepted eating practices associated with microbiome diversity, metabolic regulation, and psychosocial well-being.³² From nutritional and public health perspectives, fermented foods are accessible and culturally integrated dietary components that can serve as preventive tools.

Nutraceuticals

Nutraceuticals are notable components within the One Health framework, linking nutrition, food safety, and disease prevention across human, animal, and environmental systems. Defined as bioactive compounds derived from food sources that provide health benefits beyond basic nutrition, nutraceuticals offer targeted functional effects that can modulate host physiology and microbial dynamics. In the context of healthy nutrition, they represent a strategic extension of diet-based interventions, enabling the delivery of specific molecules, such as bioactive peptides, polyphenols, and organic acids, that influence key biological processes involved in host defense and metabolic regulation. Nutraceuticals help reduce reliance on conventional antimicrobials in both human and animal systems, thereby mitigating the emergence and spread of antimicrobial resistance.³³ These compounds can strengthen intestinal barrier function,³⁴ modulate inflammatory pathways,³⁵ and support microbiota balance,³⁶ all of which are critical determinants of susceptibility to foodborne infections. Importantly, nutraceuticals also align with the increasing demand for sustainable and preventive health strategies. Unlike pharmaceutical interventions, they can be integrated into daily diets or functional foods, offering a safe and scalable approach to improving health outcomes. Their stability, especially in the form of purified bioactive compounds or postbiotics, further enhances their applicability in food systems and clinical settings.

Fermented foods act as natural delivery vehicles for nutraceutical compounds, providing them in a structurally complex and bioavailable form. Unlike isolated supplements, food matrices facilitate synergistic interactions between microbial metabolites and host systems, enhancing their stability and functional efficacy within the gastrointestinal environment.³⁷ In addition, fermentation can reduce antinutritional factors, such as phytates, tannins, lectins, and oxalates,³⁸ and improve nutrient digestibility, further supporting the bioactivity of nutraceutical components.³⁹

Nonviable Microbial Products

The positive effects of probiotics are not restricted to live microorganisms; nonviable microbial products and dietary bioactives can also exert similar immunomodulatory and protective functions, supporting the growing interest in nutraceuticals as stable and targeted preventive agents. Postbiotics, paraprobiotics, also called parabiotics, and cell-free supernatants (CFSs) have emerged as mechanistically defined functional agents that extend the biological relevance of fermented foods beyond the requirement for viable microorganisms.⁴⁰ Postbiotics are broadly defined as preparations of inactive microorganisms and their metabolites

that confer a health benefit to the host, encompassing metabolites, cell wall fragments, extracellular vesicles, and soluble bioactive molecules derived from microbial activity. Compared with live probiotics, postbiotics offer several advantages, including enhanced safety profiles, improved physicochemical stability, and reproducibility in formulation, particularly in complex food matrices and clinical settings where viability and colonization are not guaranteed.⁴¹ Paraprobiotics, often called “ghost probiotics” or inactivated probiotics, are nonviable microbial cells or fractions, such as cell wall components, that provide health benefits to the host,⁴² while CFSs can be defined as liquid, cell-free metabolites obtained by centrifuging and filtering microbial cultures, often probiotic bacteria such as lactic acid bacteria. These attributes position nonviable microbial products as highly controllable and scalable interventions within both food systems and host-directed therapeutic strategies. At the mechanistic level, these products exert multifaceted biological effects that target both microbial pathogens and host cellular responses. Representative mechanistic effects of nonviable microbial products on epithelial integrity, inflammatory signaling, and pathogen suppression are summarized in Table 2.^{43–54}

CHALLENGES IN THE IMPLEMENTATION OF FUNCTIONAL FOOD COMPONENTS

Despite growing recognition of functional food components within the One Health framework, several critical challenges limit their consistent application and impact on health. These limitations include product variability, safety considerations, and insufficient mechanistic and clinical validation, all of which must be addressed to enable their integration into evidence-based food safety and public health strategies. The absence of standardized protocols for the characterization, quantification, and quality control of bioactive components, such as postbiotics and parabiotics, limits the ability to ensure consistency and efficacy. Challenges in balancing innovation with safety and scientific substantiation complicate the evaluation of the complex interactions between the functional properties of multicomponent food matrices and pathogens.⁵⁵

One of the primary challenges associated with fermented foods is the inherent variability in their composition and functional properties. Fermentation is a biologically dynamic process influenced by raw material quality, starter cultures, processing conditions, and environmental factors. As a result, the concentration and profile of bioactive compounds, including organic acids, peptides, and antimicrobial metabolites, can vary significantly across batches and production systems. This variability is particularly pronounced in traditional fermented ASFs, in which spontaneous fermentation and undefined microbial consortia are common. Consequently, the

Table 2. Positive effects of selected nonviable microbial products on health

Nonviable microbial product	Source microorganism	Health effects	Ref
Postbiotic	<i>Lactiplantibacillus plantarum</i>	Enhancement of tight junction proteins, including ZO-1, occludin, and claudin-1	43
Postbiotics	LAB strains	Reduction of apoptosis and oxidative stress in infected epithelial cells	44
Postbiotics	<i>Bacillus amyloliquefaciens</i> J and <i>L. plantarum</i>	Antimicrobial activity against <i>Salmonella Typhimurium</i> , <i>Staphylococcus aureus</i> , and <i>Escherichia coli</i> O157:H7	45
Postbiotic	<i>L. plantarum</i>	Reduction of bacterial invasion of epithelial cells	46
Postbiotics	<i>Bifidobacterium animalis</i> subsp. <i>lactis</i> , <i>L. plantarum</i> , and <i>Lactobacillus paracasei</i>	Suppression of the TLR4/NLRP3/MyD88 signaling pathway	47
Postbiotic	<i>Lactobacillus helveticus</i>	Improvement in stool consistency and relief of constipation-related abdominal discomfort	48
Paraprobiotics	<i>Lactobacillus casei</i> , <i>L. paracasei</i> , <i>Lactobacillus gasseri</i> , <i>B. lactis</i> , and <i>Streptococcus thermophilus</i>	Anti-inflammatory effects and reductions in organ nitrite/nitrate levels	49
Paraprobiotics	<i>L. plantarum</i> and <i>Limosilactobacillus fermentum</i>	Inhibition of the NF-κB/NLRP3 signaling pathway	50
Paraprobiotics	<i>L. plantarum</i> , <i>Lactobacillus sakei</i> , and <i>Limosilactobacillus reuteri</i>	Enhanced antioxidant activity, anti-inflammatory effects, and barrier integrity	51
Cell-free supernatants	<i>L. plantarum</i> , <i>L. paracasei</i> , <i>L. gasseri</i> , and <i>S. thermophilus</i>	Modulation of mucus production	52
Cell-free supernatants	<i>Lactobacillus amylovorus</i>	Modulation of TLR4-independent inflammatory signaling	53
Cell-free supernatants	<i>L. reuteri</i> , <i>Lactocaseibacillus rhamnosus</i> , and <i>L. plantarum</i>	Modulation of inflammation and oxidative stress	54

LAB: Lactic acid bacteria; ZO: Zonula occludens; TLR: Toll-like receptor; NLRP3: NLR family pyrin domain containing 3; MyD88: Myeloid differentiation primary response 88; NF-κB: Nuclear factor kappa-light-chain-enhancer of activated B cells; Ref: Reference.

reproducibility of functional effects, including antimicrobial activity and host-modulatory capacity, remains difficult to standardize, limiting comparability across studies and applications.⁵⁶

Although fermentation is traditionally associated with improved safety and preservation, it can also introduce specific risks if not properly controlled. The formation of biogenic amines, such as histamine, tyramine, and putrescine, through microbial decarboxylation of amino acids represents a significant safety concern, particularly in protein-rich fermented animal products.⁵⁷ Elevated levels of these compounds can lead to adverse health effects, including allergic and hypertensive reactions in sensitive individuals. In addition, inadequate hygiene or uncontrolled fermentation conditions may allow the survival or proliferation of undesirable saprophytes and mycotoxigenic molds. These

risks underscore the importance of controlled fermentation processes, validated starter cultures, and rigorous monitoring systems to ensure product safety.⁵⁸

High salt content is a notable limitation of many fermented foods, particularly animal-sourced products such as cheeses and cured or fermented meats, in which sodium chloride is essential for microbial control, texture development, and shelf-life extension. Although salt contributes to the safety and stability of these products by inhibiting undesirable microorganisms and shaping fermentation dynamics, excessive dietary sodium intake is strongly associated with adverse health outcomes, including hypertension and increased cardiovascular risk. This presents a critical paradox in the context of functional nutrition and One Health, in which foods intended to confer health benefits may simultaneously contribute to the burden of noncommunicable diseases if

consumed inappropriately. Furthermore, efforts to reduce salt content in fermented foods must be carefully balanced, as lowering sodium levels can alter microbial ecology and potentially compromise product quality. Therefore, optimizing salt concentrations, developing alternative preservation strategies, and engineering low-sodium fermentation systems remain essential research priorities to ensure the health-promoting potential of fermented foods.⁵⁹

Nutraceuticals face several key limitations that constrain their broader application and clinical translation, including a lack of standardization, limited bioavailability, regulatory ambiguity, and insufficient mechanistic and clinical evidence. Variability in composition and active compound concentrations reduces reproducibility and complicates dose optimization, while many bioactive molecules exhibit poor absorption or rapid metabolism, limiting their *in vivo* efficacy. In addition, inconsistent regulatory frameworks across regions create challenges in quality control, safety assessment, and validation of health claims. Although nutraceuticals show promising antimicrobial, antioxidant, and immunomodulatory effects against pathogens in experimental models, robust human clinical data and detailed mechanistic insights remain limited. Furthermore, potential safety concerns and interactions, particularly at higher doses or in sensitive populations, underscore the need for rigorous evaluation.⁶⁰

Despite promising *in vitro* and preclinical findings, a substantial gap remains in mechanistic clarity and clinical validation regarding the health effects of functional food compounds. Many studies rely on experimental models that do not fully capture the complexity of host–microbiota–pathogen interactions *in vivo*. Although evidence supports the antimicrobial and immunomodulatory effects of fermentation-derived compounds against pathogens, the precise molecular mechanisms and dose–response relationships remain incompletely elucidated. Moreover, well-designed clinical trials assessing efficacy, safety, and long-term outcomes in diverse populations are limited. This lack of robust clinical evidence restricts the ability to formulate definitive dietary recommendations and hinders acceptance within clinical and regulatory frameworks.

CONCLUSION

Fermented ASFs and their functional components are multifaceted, biologically active tools within the One Health framework, bridging the interconnected domains of food safety, nutrition, and host defense. Beyond their traditional roles in preservation and nutrient delivery, these functional components can exert targeted antimicrobial effects against major foodborne pathogens while simultaneously modulating intestinal barrier integrity and innate immune responses.

This dual functionality, controlling pathogens within food matrices and enhancing host resilience after consumption, positions them as strategic interventions capable of addressing both exposure risk and disease susceptibility. Importantly, integrating fermented ASFs into dietary practices, alongside nutraceuticals, postbiotics, and other bioactive metabolites, offers a complementary approach to conventional food safety systems. These strategies align with the increasing need to reduce reliance on antimicrobials, mitigate antimicrobial resistance, and implement sustainable preventive health measures across populations. Their capacity to modulate gut microbiota composition, regulate inflammatory signaling pathways, and reinforce epithelial defenses highlights their relevance in reducing the burden of foodborne diseases. From a translational perspective, the strategic incorporation of these functional approaches into food production systems, animal feeding practices, and public health policies holds significant promise. However, their successful implementation requires continued efforts toward standardization, mechanistic validation, and clinical substantiation. Within a holistic One Health context, fermented ASFs and nutraceuticals, when used as integrated tools, offer a forward-looking pathway to enhance food safety, strengthen host defense mechanisms, and promote sustainable health across interconnected biological systems.

Conflict of Interest: The author have no conflicts of interest to declare.

Funding: The author declared that this study received no financial support.

Use of AI for Writing Assistance: The author acknowledges the use of large language model–based tools, such as Grammarly, provided by Erciyes University Dean of Research to assist with English grammar, language refinement, and overall fluency during the preparation of this review. The author retains full responsibility for the scientific content, accuracy, interpretations, and conclusions presented in this review.

Peer-review: Externally peer-reviewed.

REFERENCES

1. Sheffield S, Fiorotto ML, Davis TA. Nutritional importance of animal-sourced foods in a healthy diet. *Front Nutr* 2024;11:1424912. [CrossRef]
2. Heredia N, García S. Animals as sources of food-borne pathogens: A review. *Anim Nutr* 2018;4(3):250-5. [CrossRef]
3. Park I, Mannaa M. Fermented Foods as Functional Systems: Microbial Communities and Metabolites Influencing Gut Health and Systemic Outcomes. *Foods* 2025;14(13):2292. [CrossRef]
4. Soemarie YB, Milanda T, Barliana MI. Fermented Foods as Probiotics: A Review. *J Adv Pharm Technol Res* 2021;12(4):335-9. [CrossRef]

5. van Zyl WF, Deane SM, Dicks LMT. Molecular insights into probiotic mechanisms of action employed against intestinal pathogenic bacteria. *Gut Microbes* 2020;12(1):1831339. [\[CrossRef\]](#)
6. Zimmermann C, Wagner AE. Impact of Food-Derived Bioactive Compounds on Intestinal Immunity. *Biomolecules* 2021;11(12):1901. [\[CrossRef\]](#)
7. World Health Organization. Strengthening surveillance of and response to foodborne diseases. Stage three manual: integrating surveillance data to better understand risks across the food chain. <https://www.who.int/publications/i/item/9789240118201> Accessed June 11, 2026.
8. World Health Organization (WHO). Estimating the burden of foodborne diseases. 2026. <https://www.who.int/activities/estimating-the-burden-of-foodborne-diseases> Accessed June 11, 2026.
9. USDA, Economic Research Service (ERS). Cost Estimates of Foodborne Illnesses. <https://www.ers.usda.gov/data-products/cost-estimates-of-foodborne-illnesses> Accessed March, 2026.
10. Tong X, Zhang H. Perception, Amplification and Communication: A Case Study of Food Safety Risks. *China's Emergency Management* 2019:71-115. [\[CrossRef\]](#)
11. Richardson LC, Cole D, Hoekstra RM, Rajasingham A, Johnson SD, Bruce BB. Foods Implicated in U.S. Outbreaks Differ from the Types Most Commonly Consumed. *J Food Prot* 2021;84(5):869-75. [\[CrossRef\]](#)
12. Oltean HN, Lipton B, Black A, Snekvik K, Haman K, Buswell M, et al. Developing a one health data integration framework focused on real-time pathogen surveillance and applied genomic epidemiology. *One Health Outlook* 2025;7(1):9. [\[CrossRef\]](#)
13. Shrivastava P, Sahoo SK, Venkatesh KM. Role of veterinarians in public health. *Epidemiology and environmental hygiene in veterinary public health*. John Wiley & Sons; 2025.p.287-298. [\[CrossRef\]](#)
14. Bold J, Rostami K. Foodborne, food related illness and role of the healthcare professionals. *Gastroenterol Hepatol Bed Bench* 2011;4(1):1-2.
15. Arredondo-Rivera M, Barois Z, Monti GE, Steketee J, Daburon A. Bridging Food Systems and One Health: A key to preventing future pandemics? *One Health* 2024;18:100727. [\[CrossRef\]](#)
16. Zhang D, Jian YP, Zhang YN, Li Y, Gu LT, Sun HH, et al. Short-chain fatty acids in diseases. *Cell Commun Signal* 2023;21(1):212. [\[CrossRef\]](#)
17. Wang Y, Wu J, Lv M, Shao Z, Hungwe M, Wang J, et al. Metabolism Characteristics of Lactic Acid Bacteria and the Expanding Applications in Food Industry. *Front Bioeng Biotechnol* 2021;9:612285. [\[CrossRef\]](#)
18. Al S, Franzen AJ, Hwang K, Campbell RE, Glass K, Vogel KD, et al. Comparison of peracetic acid and bacteriophage application by vascular rinsing on Salmonella reduction in lymph nodes of goat carcasses. *J Food Sci* 2025;90(4):e70203. [\[CrossRef\]](#)
19. Donkor ON, Henriksson A, Singh TK, Vasiljevic T, Shah NP. ACE-inhibitory activity of probiotic yoghurt. *Int Dairy J* 2007;17(11):1321-31. [\[CrossRef\]](#)
20. Ni WW, Zhang QM, Zhang X, Li Y, Yu SS, Wu HY, et al. Modulation effect of *Lactobacillus acidophilus* KLDS 1.0738 on gut microbiota and TLR4 expression in β -lactoglobulin-induced allergic mice model. *Allergol Immunopathol (Madr)* 2020;48(2):149-57. [\[CrossRef\]](#)
21. Martínez-Maqueda D, Miralles B, Cruz-Huerta E, Recio I. Casein hydrolysate and derived peptides stimulate mucin secretion and gene expression in human intestinal cells. *Int Dairy J* 2013;32(1):13-19. [\[CrossRef\]](#)
22. Norris GH, Jiang C, Ryan J, Porter CM, Blesso CN. Milk sphingomyelin improves lipid metabolism and alters gut microbiota in high fat diet-fed mice. *J Nutr Biochem* 2016;30:93-101. [\[CrossRef\]](#)
23. Takayama H, Fukatsu K, Takahashi K, Noguchi M, Watkins A, Matsumoto N, et al. Influences of a fermented milk with *Lactobacillus bulgaricus* and *Streptococcus thermophilus* on gut associated lymphoid tissue, mucosal IgA, and gut flora in mice. *Clin Nutr Open Sci* 2022;42:36-48. [\[CrossRef\]](#)
24. Bengoa AA, Dardis C, Gagliarini N, Garrote GL, Abraham AG. Exopolysaccharides From *Lactobacillus paracasei* Isolated From Kefir as Potential Bioactive Compounds for Microbiota Modulation. *Front Microbiol* 2020;11:583254. [\[CrossRef\]](#)
25. Chen Y, Wang Z, Chen X, Liu Y, Zhang H, Sun T. Identification of angiotensin I-converting enzyme inhibitory peptides from koumiss, a traditional fermented mare's milk. *J Dairy Sci* 2010;93(3):884-92. [\[CrossRef\]](#)
26. Qian B, Xing M, Cui L, Deng Y, Xu Y, Huang M, et al. Antioxidant, antihypertensive, and immunomodulatory activities of peptide fractions from fermented skim milk with *Lactobacillus delbrueckii* ssp. *bulgaricus* LB340. *J Dairy Res* 2011;78(1):72-9. [\[CrossRef\]](#)
27. Correias NH, Martínez AR, Abellán A, Sánchez HP, Tejada L. Curing strategies and bioactive peptide generation in ham: *In vitro* digestion and *in silico* evaluation. *Food Chem* 2025;484:144360. [\[CrossRef\]](#)
28. Phong HX, Viet LQ, Chau LM, Long BHD, Thanh NN, Phat DT, et al. Isolation and selection of lactic acid bacteria with

- the capacity of producing γ -aminobutyric acid (GABA) and antimicrobial activity: Its application in fermented meat product. *Curr Nutr Food Sci* 2023;19(8):831-7. [\[CrossRef\]](#)
29. Pérez-Burillo S, Mehta T, Pastoriza S, Kramer DI, Paliy O, Rufián-Henares JA. Potential probiotic salami with dietary fiber modulates antioxidant capacity, short chain fatty acid production and gut microbiota community structure. *LWT* 2019;105:355-62. [\[CrossRef\]](#)
 30. Song MY, Van-Ba H, Park WS, Yoo JY, Kang HB, Kim JH, et al. Quality Characteristics of Functional Fermented Sausages Added with Encapsulated Probiotic *Bifidobacterium longum* KACC 91563. *Korean J Food Sci Anim Resour* 2018;38(5):981-94. [\[CrossRef\]](#)
 31. Chin SF, Liu W, Storkson JM, Ha YL, Pariza MW. Dietary sources of conjugated dienoic isomers of linoleic acid, a newly recognized class of anticarcinogens. *J Food Compos Anal* 1992;5(3):185-97. Japanese. [\[CrossRef\]](#)
 32. Dunn RR, Wilson J, Nichols LM, Gavin MC. Toward a global ecology of fermented foods. *Curr Anthropol* 2021;62(S24):S220-32. [\[CrossRef\]](#)
 33. Ballou MA, Davis EM, Kasl BA. Nutraceuticals: An Alternative Strategy for the Use of Antimicrobials. *Vet Clin North Am Food Anim Pract* 2019;35(3):507-34. [\[CrossRef\]](#)
 34. McCarty MF, Lerner A. Perspective: Prospects for Nutraceutical Support of Intestinal Barrier Function. *Adv Nutr* 2021;12(2):316-24. [\[CrossRef\]](#)
 35. Huang MT, Ghai G, Ho CT. Inflammatory Process and Molecular Targets for Antiinflammatory Nutraceuticals. *Compr Rev Food Sci Food Saf* 2004;3(4):127-39. [\[CrossRef\]](#)
 36. Gao S, He Y, Zhang L, Liu L, Qu C, Zheng Z, et al. Conjugated linoleic acid ameliorates hepatic steatosis by modulating intestinal permeability and gut microbiota in ob/ob mice. *Food Nutr Res* 2022;66. [\[CrossRef\]](#)
 37. Wang J, Wu P, Chen XD, Yu A, Dhital S. Effect of Food Matrix and Administration Timing on the Survival of *Lactobacillus rhamnosus* GG During *in Vitro* Gastrointestinal Digestion. *Foods* 2025;14(17):3076. [\[CrossRef\]](#)
 38. Naseem A, Akhtar S, Ismail T, Qamar M, Sattar DE, Saeed W, et al. Effect of Growth Stages and Lactic Acid Fermentation on Anti-Nutrients and Nutritional Attributes of Spinach (*Spinacia oleracea*). *Microorganisms* 2023;11(9):2343. [\[CrossRef\]](#)
 39. Shahni K, Haobam B, Singh OI, Devi KS, Devi ST, Sharma N, et al. Nutraceuticals Potential of Fermented Foods: A Global and Ethnic Perspective on Health and Wellness. *Fermentation* 2026;12(5):211. [\[CrossRef\]](#)
 40. Nataraj BH, Ali SA, Behare PV, Yadav H. Postbiotics-parabiotics: the new horizons in microbial biotherapy and functional foods. *Microb Cell Fact* 2020;19(1):168. [\[CrossRef\]](#)
 41. Vinderola G, Sanders ME, Salminen S. The Concept of Postbiotics. *Foods* 2022;11(8):1077. [\[CrossRef\]](#)
 42. Lee NK, Park YS, Kang DK, Paik HD. Paraprobiotics: definition, manufacturing methods, and functionality. *Food Sci Biotechnol* 2023;32(14):1981-91. [\[CrossRef\]](#)
 43. Zhang X, Li Y, Zhang C, Chi H, Liu C, Li A, et al. Postbiotics derived from *Lactobacillus plantarum* 1.0386 ameliorate lipopolysaccharide-induced tight junction injury via MicroRNA-200c-3p mediated activation of the MLCK-MLC pathway in Caco-2 cells. *Food Funct* 2022;13(21):11008-20. [\[CrossRef\]](#)
 44. Nowak A, Zakłós-Szyda M, Rosicka-Kaczmarek J, Motyl I. Anticancer Potential of Post-Fermentation Media and Cell Extracts of Probiotic Strains: An *in Vitro* Study. *Cancers (Basel)* 2022;14(7):1853. [\[CrossRef\]](#)
 45. Tong Y, Abbas Z, Zhang J, Wang J, Zhou Y, Si D, et al. Antimicrobial activity and mechanism of novel postbiotics against foodborne pathogens. *LWT* 2025;217:117464. [\[CrossRef\]](#)
 46. Çobur H, Löker N, Dışhan A, Vurmaz M, Karatepe P, Hızlısoy H, et al. Lactiplantibacillus Plantarum Postbiotics Suppress Salmonella Typhimurium Invasion and Modulate Innate Responses in Human Intestinal Epithelial Cells. *Probiotics Antimicrob Proteins* 2026;18(3):4969-82. [\[CrossRef\]](#)
 47. Jia L, Xiao L, Song X, Zhang X, Han X, Liang W, et al. Postbiotics alleviate cognitive impairment and neuroinflammation via gut-brain axis and TLR4/MyD88/NLRP3 pathway. *Food Res Int* 2025;222(Pt 2):117708. [\[CrossRef\]](#)
 48. Tanihiro R, Yuki M, Sakano K, Sasai M, Sawada D, Ebihara S, et al. Effects of Heat-Treated *Lactobacillus helveticus* CP790-Fermented Milk on Gastrointestinal Health in Healthy Adults: A Randomized Double-Blind Placebo-Controlled Trial. *Nutrients* 2024;16(14):2191. [\[CrossRef\]](#)
 49. Michels M, Jesus GFA, Córneo E, Rocha LB, Dias R, Voytena APL, et al. Paraprobiotics Mitigate Inflammation and Anxiety-Like Behavior in a Lipopolysaccharide-Induced Mouse Model. *Curr Microbiol* 2025;82(9):385. [\[CrossRef\]](#)
 50. Liu C, Qi X, Guan K, Chen H, Li Q, Mao K, et al. Paraprobiotics 5-PA and 6-PA restore intestinal homeostasis by inhibiting NF- κ B/NLRP3 signaling pathway and alleviating dysbiosis in mice with chronic ulcerative colitis. *J Funct Foods* 2024;113:106048. [\[CrossRef\]](#)
 51. Kang YY, Song HJ, Park SY, Oh DN, Kim GY, Been NY, et al. Comparative Effects of Probiotics and Paraprobiotics Derived from *Lactiplantibacillus plantarum*, *Lactilactobacillus sakei*, and *Limosilactobacillus reuteri* in a

- DSS-Induced Ulcerative Colitis Mouse Model. *J Microbiol Biotechnol* 2025;35:e2411045. [\[CrossRef\]](#)
52. Lee JY, Kang CH. Cell-Free Supernatant from *Lactobacillus* and *Streptococcus* Strains Modulate Mucus Production via Nf- κ B/CREB Pathway in Diesel Particle Matter-Stimulated NCI-H292 Airway Epithelial Cells. *Molecules* 2022;28(1):61. [\[CrossRef\]](#)
53. Finamore A, Roselli M, Imbinto A, Seeboth J, Oswald IP, Mengheri E. *Lactobacillus amylovorus* inhibits the TLR4 inflammatory signaling triggered by enterotoxigenic *Escherichia coli* via modulation of the negative regulators and involvement of TLR2 in intestinal Caco-2 cells and pig explants. *PLoS One* 2014;9(4):e94891. [\[CrossRef\]](#)
54. Di Chiano M, Rocchetti MT, Spano G, Russo P, Allegretta C, Milior G, et al. *Lactobacilli* Cell-Free Supernatants Modulate Inflammation and Oxidative Stress in Human Microglia via NRF2-SOD1 Signaling. *Cell Mol Neurobiol* 2024;44(1):60. [\[CrossRef\]](#)
55. Divsalar E, Incili GK, Shi C, Semsari E, Hosseini SH, Ebrahimi Tirtashi F, et al. Challenges with the use of postbiotics/parabiotics in food industry. *Crit Rev Food Sci Nutr* 2026;66(6):1155-71. [\[CrossRef\]](#)
56. Pérez-Burillo S, Molino S, Navajas-Porras B, Valverde-Moya ÁJ, Hinojosa-Nogueira D, López-Maldonado A, et al. An *in vitro* batch fermentation protocol for studying the contribution of food to gut microbiota composition and functionality. *Nat Protoc* 2021;16(7):3186-209. [\[CrossRef\]](#)
57. Saha Turna N, Chung R, McIntyre L. A review of biogenic amines in fermented foods: Occurrence and health effects. *Heliyon* 2024;10(2):e24501. [\[CrossRef\]](#)
58. Skowron K, Budzyńska A, Grudlewska-Buda K, Wiktorczyk-Kapischke N, Andrzejewska M, Wałęcka-Zacharska E, et al. Two Faces of Fermented Foods-The Benefits and Threats of Its Consumption. *Front Microbiol* 2022;13:845166. [\[CrossRef\]](#)
59. Lin X, Tang Y, Hu Y, Lu Y, Sun Q, Lv Y, et al. Sodium Reduction in Traditional Fermented Foods: Challenges, Strategies, and Perspectives. *J Agric Food Chem* 2021;69(29):8065-80. [\[CrossRef\]](#)
60. Rajalakshmi R, Melians MA, Pon FF, Cosio DS, Buvaramurthy V, Jayakumar AR, et al. Advantages and disadvantages of nutraceuticals. *Nutraceuticals for Alzheimer's Disease: A Promising Therapeutic Approach*. Springer Nature Singapore;2023.p.245-286. [\[CrossRef\]](#)