

Characterization of an Egyptian *Spodoptera littoralis* Nucleopolyhedrovirus with Cloning and Sequencing of a Highly Conserved Region From Polyhedrin Gene

Polihedrin gende iyi korunmuş bölgenin klonlanması ve zincirlenmesi ile bir Mısır *Spodoptera littoralis* Nükleopolihedrovirüs'ün karakteristiği

AlaaEddeen Seufi,

Assoc. Prof.,
Department of Entomology,
Faculty of Science, Cairo University,
Giza, Egypt.
alaaseufi@yahoo.com

Abstract

Purpose: An Egyptian isolate of *Spodoptera littoralis* nucleopolyhedrovirus (SpliNPV) was tested for its potential as biocontrol agent in comparison to *Autographa californica* multiple nucleopolyhedrovirus (AcMNPV).

Material and Methods: Comparative assays of SpliNPV and AcMNPV against 2nd instar larvae of *Spodoptera littoralis* revealed 4-fold greater susceptibility of *S. littoralis* to AcMNPV than to SpliNPV based on LC₅₀ values for the two viruses. The LT₅₀s determined for SpliNPV and AcMNPV using LC₅₀ of the virus against 2nd instar larvae were 4.2 and 5.8 days, respectively. A DNA segment of 405 bp containing highly conserved region from polyhedrin gene of SpliNPV (Polh-cr) was successfully amplified by PCR. Subsequently, this DNA segment was cloned and sequenced. Nucleotide sequence and its deduced amino acid sequence were compared to all available sequences in GenBank.

Results: Sequence alignment results revealed that Polh-cr showed significant similarities with 91 different baculovirus isolates. The percentage of homology ranged from 78% for *Plusia orichalcea* NPV to 99 % for SpliNPV.

Conclusion: This highly conserved region provides a candidate that could be used in easy, fast and economic prospective systems for virus detection as well as in biological control strategies.

Keywords: **Polyhedrin; Nucleopolyhedrovirus ; Spodoptera; Polimerase chain reaction.**

This manuscript can be downloaded from the webpage:
[http://tjpdergisi.erciyes.edu.tr/project6/2007;29\(5\)353-362.pdf](http://tjpdergisi.erciyes.edu.tr/project6/2007;29(5)353-362.pdf)

Submitted : August 19, 2006
Revised : June 27, 2007
Accepted : July 17, 2007

Corresponding Author:

A.M.Seufi,
Department of Entomology,
Faculty of Science, Cairo University,
Giza, Egypt.

Telephone : +20 - 202-7010367
E-mail : alaaseufi@yahoo.com

Özet

Amaç: Bir Mısır *Spodoptera littoralis* Nucleopolyhedrovirus (SpliNPV) potansiyel biokontrol ajanı olabilmesi için *Autographa californica* (AcMNPV) ile karşılaştırılarak test edildi.

Material and Methods: *Spodoptera littoralis* ikinci dönem larvasına karşı SpliNPV ve AcMNPV virüsünün LC₅₀ değerine dayanan karşılaştırmalı değerlendirmede AcMNPV'nin SpliNPV göre dört kat daha büyük duyarlılık gösterdiği ortaya çıkarılmıştır. *Spodoptera littoralis* ikinci dönem larvasına karşı SpliNPV ve AcMNPV virüsünün yaklaşık LC₅₀ değeri kullanılarak belirlenen LT₅₀ sırasıyla 4.2 ve 5.8 gündür. PCR, SpliNPV'nin polyhedrin geninin iyi korunmuş bölgesini başarıyla amlifiye etti.

405 bp'lık bir DNA geni (Polh-cr) elde edildi. Bundan sonra bu DNA segmenti klonlandı ve zincirlendi. Nükleotid dizilimi ve amino asit dizilimi GenBank'ta bütün olabilen dizimlerle karşılaştırıldı.

Bulgular: Polh-cr'den üretilen önemli dizim sonuçları 91 Baculovirus izolatları ile ilişkilendirildi. Homoloji oranları *Plusia orichalcea* NPV için %78 ve SpliNPV için %99 olarak bulundu.

Sonuç: Biyolojik kontrol stratejilerinde virüs belirlemek için bu yüksek olarak korunmuş bölgenin kullanılması muhtemel kolay, hızlı ve ekonomik olarak görülmüyor.

Anahtar kelimeler: **Nükleopolihedrovirüs; Polimeraz zincir reaksiyonu; Polihedrin; Spodoptera.**

Introduction

Baculoviruses are considered to be the largest and most broadly studied insect viruses. Although they infect over 600 species of insects (1), individual isolates normally show a limited host range and infect only closely related species. It is believed that baculoviruses are potentially useful as safe biological control agent and in some cases they were used successfully to control different insect pests (2,3,4). However the overall use of baculoviruses for biological control is limited compared to other pest control means (3,5). This is due to the virulence and speed of action, as related to dose, host range, cost of production and patent registration are important effectiveness-determining properties of insect-pathogenic biocontrol agents.

Baculoviruses were not only isolated from the insect orders Hymenoptera, Diptera and Trichoptera, but also were isolated from the crustacean order Decapoda (shrimps) (6). Furthermore, some NPVs (*Penaeus monodon*-NPV) are considered as serious pests for marine crustaceans (7). The infection of marine crustaceans (shrimps, prawns, crabs, eustacos, etc.) with NPV will reduce their economic value and thus negatively affecting the national income. This problem will arise in countries that depend on marine wealth as a source of national income and/ or where marine crustaceans are a major in most of the peoples' food (Australia and Far East nations like Japan, China, etc.) For the above mentioned reasons, many authors were interested in developing an accurate and easy diagnostic method to detect baculoviral infection in larval shrimps as well as insects (7-18). Because baculoviruses are of great interest and utility to a large cross-sections of agricultural and biomedical research community, selection of some isolates with natural improved characteristics and definitive characterization of them have appeared recently. Consequently this piece of work will help in viral epidemiology, in monitoring the viral dissemination after field application and in risk assessment studies. Therefore, the main objective of the present work is to obtain a sequenced highly conserved DNA fragment from our NPV isolate to be used in an easy, fast and economic prospective system for virus detection. Also, this DNA segment can be used for developing kits of ELISA, hybridization, Western and dot blotting. Furthermore, our NPV isolate may be useful as potential biocontrol agent in programs of integrated pest management (IPM).

Materials and Methods

Insects. Laboratory colony of the cotton leaf worm, *Spodoptera littoralis*, was originally collected from Giza, Egypt and maintained in the insectary of Agricultural Genetic Engineering Research Institute (AGERI) under highly controlled conditions from 1990 to date. The colony was maintained in the laboratory according to the technique described by El-Defrawi et al., (19). Larvae were reared on a semisynthetic diet described by Levinson and Navon (20). This colony was kept at 25 (23-27)°C, 65-70% RH and natural photoperiod. These insects were used for viral propagation and bioassays.

Viruses. An NPV isolate was selected from thirteen Egyptian baculovirus isolates from different geographical localities during 1996-2000. This isolate was studied in detail to determine its insecticidal activity compared to AcMNPV. In addition, a highly conserved region of polyhedrin gene was amplified, cloned and sequenced for further investigations. The selected isolate was collected from an Egyptian field located at Khorshid, Alexandria. It was isolated during Jan, 1999 from a *S. littoralis* larva on cabbage plants. This isolate was successfully propagated, purified following the method described by Lacey et al. (21) and used for further studies.

A virus stock of NPV from *Autographa californica* nuclear polyhedrosis virus (AcMNPV; E2 strain) was originally obtained from Prof. Dr. Suzanne Thiem, Department of Entomology, University of Michigan, was used in the bioassays as a reference in comparison to our isolate.

Insecticidal activity. Laboratory bioassay tests were conducted using highly purified virus suspensions. The SpliNPV and AcMNPV isolates were tested against 2nd larval instar from the virus-free rearing culture in the laboratory. The concentrations were measured using Thoma haemocytometer and light microscope (21). Bioassays were performed using five concentrations of each virus isolate. Tested larvae were starved for 8 hours prior to feeding viruses. Serial dilutions of each tested isolate were prepared beginning with the following stock concentrations in PIB/ ml: 7.8×10^4 and 9.4×10^4 for SpliNPV and AcMNPV, respectively. Bioassays were carried out in plastic cups measuring 15 cm diameter x 10 cm height and containing a layer of 1 cm semi-synthetic diet. The viral suspensions were dispersed on the diet using micropipette (25 µl suspension/ larva). Forty larvae were used for each concentration of virus and control. The control larvae were fed on diet treated with distilled

water. All treatments were kept at 25 (23-27)°C and normal photoperiod. The experiment was replicated thrice for each virus isolate and control. Mortality was recorded on the 7th day post-infection then corrected according to Abbott's formula (22). Cumulative mortality was recorded daily and the experiment was stopped on day 10 post-infection. Mortality was corrected according to Abbott's formula (22).

Statistical Analyses. Probit analysis of mortality data from bioassays was conducted using SPSS (ver 10.0) computer software (SPSS for Windows, SPSS Inc., 1997). The LT values for the tested isolates were derived from analysis of data on the progression of mortality of *S. littoralis* larvae, following exposure to the LC₅₀s dosage using probit analysis modified for multiple observations over time (Throne et al., 23) and Mathematica software (Wolfram, Champaign, IL).

PCR amplification. PCR amplification was performed according to Saiki et al. (24) with minor modifications. Total DNA was extracted from the NPV isolate and the DNA segment was amplified using two primers designed based on conserved nucleotide sequences of ten different polyhedrin genes (Abdel-Hamid et al., 25). The forward primer: GG(GT) CC(GT) GGC AAG AAT CAG AA and the reverse one: GCG TC(TG) GG(TG) GCG AAC TCT TT(TG) ATT TT. Total reaction volume was 50 µl which contained 1x PCR buffer (Promega), 1.5 mM MgCl₂, 200 µM dNTPs, 2.5 U Taq DNA polymerase (Promega), 100 ng of each primer and 30 ng of template DNA. The amplification program used was 3 min at 94 °C (hot start), 1 min at 94 °C, 2 min at 55 °C and 2 min at 72 °C for 35 cycles followed by one cycle of 72 °C for 7 min. PCR amplification was carried out in a DNA thermal cycler (Model 380 A, Applied Biosystems, CA, USA). Cloning of a highly conserved region from SpliNPV polyhedrin gene. The positive PCR products were visualized and eluted from the gel using GenClean Kit (Invitrogen Corporation, San Diego, CA, USA) as described by the manufacturer. The purified PCR products and a pGEM-T vector (Promega Corporation, Madison, WI, USA) were mixed in a 5: 1 (insert: vector) molar ratio and ligated using T4 DNA ligase (as described by the manufacturer). Ligation mix was used to transform competent *Escherichia coli* JM109 cells (Stratagene, La Jolla, CA, USA). White colonies were screened using PCR as described earlier in this section.

Nucleotide sequence and sequence analysis. Only one clone pGNPV-95 was selected and sequenced using M13 universal forward and reverse primers. Sequencing was performed using T⁷ SequencingT^M kit (Pharmacia, Biotech) and model 310 automated sequencer (Applied Biosystems, Foster City, CA, USA). Analysis of nucleotide and deduced amino acid sequences was carried out using EditSeq-DNAstar Inc., Expert Sequence Analysis software, Windows 32 Edit Seq 4.00 (1989-1999) and ExPasy database on the internet. Blast search for alignment of the obtained sequence with the published ones was done using database of National Centre for Biotechnology Information (NCBI). The cloned DNA fragment (Polhcr) was deposited in GenBank under the AY442260 accession number.

Results

Field survey. During the present study, the most promising isolate, as a biocontrol agent (based on LC and LT values), was selected from thirteen Egyptian baculovirus isolates. Through four visits to the location, out of 63 collected *S. littoralis* larvae, 18 larvae were diseased (showed symptoms of viral infection). The virus was diagnosed using light microscope, propagated in a *S. littoralis* laboratory colony, purified and kept at -80 °C for further studies.

Insecticidal activity. Table I presents the LC and LT values for the two tested isolates (SpliNPV and AcMNPV). It is clear that the newly molted 2nd larval instars of *S. littoralis* were susceptible to the applied concentrations. Based on LC₅₀ in PIB/ ml, our SpliNPV isolate was significantly (4-fold) less active (LC₅₀= 1.2X10⁻³) than the reference strain AcMNPV (LC₅₀= 3.7X10⁻²). Although LC₂₅ and LC₉₅ of our isolate were approximately 2 and 8-fold lower than that of AcMNPV, there were substantial overlap in the 95% confidence limits of the two viruses. Time to death showed some dependence on the initial concentration. Furthermore, it was observed that the highest peak of mortality was on 4-6 days post-infection. The progression of mortality in 2nd instar larvae of *S. littoralis* over a 10-day-period following exposure to LC₅₀ dosage of the two viruses was analyzed to produce LT values (Table I). Probit analysis of the LT values revealed that the decrease in LT values in favour of our isolate against AcMNPV was not significant. Despite the overlap in the 95% confidence intervals of the two viruses, LT₉₅ of our isolate was nearly 4 days less than that of AcMNPV (95% c.i. 9.7-25.4 days, X² = 2.64, DF = 8, P = 0.96, slope = 3.49±0.74).

PCR Amplification. Two oligonucleotide primers were designed to amplify 537 bp within the open reading frame (orf) of polyhedrin gene (25) and were successfully used in PCR. PCR analysis of our results revealed that a DNA fragment of only 405 bp was amplified within the orf (beginning about 78 - 81 codons after the starting codon, AUG) of the SpliNPV-95 polyhedrin gene (Fig. 1; panel B, lane 1). Subsequently this DNA segment was cloned into pGEM-T vector (Fig. 1; panel A, lane 2) and transformed cells were tested with PCR using the same primers (Fig. 1; panel B, lane 2). Using this screening method, clone pGNPV-95 was tested as positive control (Fig. 1; panel A, lane 2).

Nucleotide sequence and sequence analyses. The nucleotide sequence of Polh-cr and its deduced amino acid sequence are shown in Fig 2. A single open reading frame (orf) that could encode a polypeptide of 135 amino acids was detected. No stop codon was found all over the sequence. This deduced polypeptide contains 16 strongly basic, 16 strongly acidic, 46 hydrophobic and 34 polar amino acids. The calculated molecular mass of the putative polypeptide is 15.92 KDa. Isoelectric point (PI) is 7.242 and the charge at pH 7.0 is 0.331.

The nucleotide sequence of Polh-cr was blasted in GenBank database and compared to all available sequences. Alignment results revealed that Polh-cr has significant alignment with 111 baculovirus isolates (100 NPVs and 11 GVs). The percentage of homology ranged between 99 % for SpliNPV (Acc# D01017) and 78 % for Plusia orichalcea NPV (Acc# AF019882). Interestingly Polh-cr produced significant alignments with 11 granulovirus sequences. This may raise the question of possible homologous recombination between NPV and GV species.

Comparing Polh-cr nucleotide sequence (Acc# AY442260) to its corresponding sequence of AcMNPV (Acc# M25054) as a reference, 82% homology, 57 different nucleotides and 12 gaps were observed throughout the compared DNA segments (Fig 3).

The deduced amino acid sequence was compared to all other polyhedrins in GenBank database. Alignment results showed that the percentage of homology of Polh-cr ranged between 100 % for *S. littoralis* polyhedrins (Acc# AAC33752 and AAR04375) and 81 % for *Attacus ricini* polyhedrin (Acc# P31036).

Table I: Susceptibility of 2nd instar larvae of *S. littoralis* to the nucleopolyhedroviruses of *A. californica* (AcMNPV) and *S. littoralis* (SpliNPV).

Isolate	LC values in PIB/ml (95% confidence limits)			LT values in days (95% confidence limits)		
	LC ₂₅	LC ₅₀ *	LC ₉₅	LT ₂₅	LT ₅₀	LT ₉₅
AcMNPV	19.5 (4.8 – 4.0 x 10)	3.7x10 ² (1.9 x 10 ² – 6.7 x 10 ²)	7.5x10 ⁵ (2.4 x 10 ⁵ – 3.7 x 10 ⁶)	3.7 (2.50 – 4.59)	5.8 (4.73 – 7.14)	17.2 (12.01 – 39.03)
SpliNPV	39.3 (1.4 x 10 – 8.2 x 10)	1.2x10 ³ (6.9 x 10 ² – 2.3 x 10 ³)	6.1x10 ⁶ (4.5 x 10 ⁵ – 1.9 x 10 ⁷)	2.6 (1.65 – 3.39)	4.2 (3.26 – 5.17)	13.4 (9.74 – 25.42)

Significant difference at 95% confidence interval.

On comparing amino acid sequence of the putative polypeptide of Polh-cr (Acc# AAR04375) to its corresponding sequences of AcMNPV (Acc# AAA46736) and other 4 SpliNPVs (Acc# AAW49208, BAA00824, AAT10182 and AAW49207), 15 different amino acids were observed throughout the compared putative polypeptides using AcMNPV as a reference (Fig. 4). It is note worthy to mention that all SpliNPVs amino acid sequences (Acc# AAW49208, BAA00824, AAT10182 and AAW49207) were identical to Polh-cr putative polypeptide (Acc# AAR04375) except for the position 133 (I a.a. was replaced with V a.a.).




Figure 1. Agarose gel electrophoresis showing clone pGNPV-95 after linearization with NdeI, and PCR confirmation. Lanes 1 and 2 show pGEM-T and pGNPV-95 after digestion with NdeI, respectively. Lanes 3 and 4 show Lambda DNA/ HindIII Marker. Lanes 5, 6 and 7 show the 405 bp amplified DNA segment from SpliNPV (as positive control), from

```

GTGAAGCCCGATACGATGAAGCTGATCGTCAACTGGAACGGCAAAGAGTTT      CTCCGTGAG
V K P D T M K L I V N W N G K E F L R E

ACTTGGACCCGTTTCATGGAAGACAGCTTCCCCATCGTGAACGATCAAGAA      GTGATGGAC
T W T R F M E D S F P I V N D Q E V M D

GTGTTTCTAGTGGTGAACATGCGTCCCAGTACGACCGAACC GTTGTCTTTAG      ATTTTTGGCG
V F L V V N M R P T R P N R C F R F L A

CAACACGCGCTCCGATGCGACCCCGAGTACGTTCCCCACGACGTGATCCGC      ATCGTCGAA
Q H A L R C D P E Y V P H D V I R I V E

CCGTCGTACGTCGGCACCAACAATGAATACCGCATCAGTCTCGCCAAGAAA      GGTGGCGGC
P S Y V G T N N E Y R I S L A K K G G G

TGTCCCGTGATGAACCTGCACGCCGAATACCACTTCGTTTGAGAGTTTC      ATCGACAAG
C P V M N L H A E Y T T S F E S F I D K

GTGATATGGTACAAC TTTTACAAGCCCATCGTCTACATCGGCACA
V I W Y N F Y K P I V Y I G T

```

Figure 2: Nucleotide and corresponding deduced amino acid sequence of a highly conserved region of polyhedrin gene (polh-cr).

```
SpliNPV 4  AAGCCCGATACGATGAAGCTGATCGTCAACTGGAACGGCAAAGAGTTTCTCCGTGAGACT 63
||||| ||||| | ||||| ||||| ||||| | |||
AcMNPV 1484 AAGCCCGACACCATGAAGCTGGTAGTAAACTGGAGCGGCAAAGAGTTTCTCAGGGAAACT 1543

SpliNPV 64  TGGACCCGTTTTCATGGAAGACAGCTTCCCCATCGTGAACGATCAAGAAAGTATGGACGTG 123
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
AcMNPV 1544 TGGACCCGTTTTCATGGAAGACAGCTTCCCTATCGTGAACGACCAAGAAATTATGGACGTG 1603

SpliNPV 124 TTTCTAGTGGTGAACATGCGTCCCCTAGACCGAACCGTTGCTTT-AGATTTTGGCGCA 182
|| |||| | ||||| || || || |||| |||||
AcMNPV 1604 TTCCTAGTGATCAACATGAGACCCACTAGGCCAACAGATG-TTTCAGATTCTGGCGCA 1662

SpliNPV 183 ACACGCGCTCCGATGCGACCCCGAGTACGTTCCCACGACGTGATCCGCATCGTCAACC 242
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
AcMNPV 1663 ACACGCTCTCCGTTGCGACCCCTGACTACGTTCCCTACGAAGTCAATCGCATTGTGGAGCC 1722

SpliNPV 243 -GTCGTACGTCGGCACCAACAATGAATACCGCATCAGTCTCGCCAAGAAAGGTGGCGGCT 301
|| |||| | ||||| ||||| ||||| ||||| |||||
AcMNPV 1723 CGT-GTACGTAGGAAACAACAACGAATACCGCATCAGCCTGGCCAAGAAGGGCGGCGGCT 1781

SpliNPV 302 GTCCC-GTGATGAACCTGCACGCCGAATACAC-CACTTCGTTTGA-GAGTTTCATCGACA 358
|| |||| || || |||| ||||| ||||| ||||| ||||| |||||
AcMNPV 1782 G-CCCTGTCATGAACCTTCACTCTGAGTACACGCACTCC-TTCGAAGAGTT-CATCAACC 1838

SpliNPV 359 AG-GTGATATGGTACAACCTTTTACAAGCCCATCGTCTAC 396
|| || |||| ||||| ||||| |||||
AcMNPV 1839 -GTGTCATCTGGGAGAACTTCTACAAACCCATCGTCTAC 1876
```

Figure 3. Comparison of Polh-cr nucleotide sequence (Acc# AY442260) to its corresponding sequence of AcMNPV (Acc# M25054) as a reference. Gaps and different nucleotides are shaded.

```
SpliNPVa: V K P D T M K L I V N W N G K E F L R E T W T R F M E D S F P I V N D Q E V M D V F L
AcMNPV: V K P D T M K L V V N W S G K E F L R E T W T R F M E D S F P I V N D Q E I M D V F L
SpliNPV1: V K P D T M K L I V N W N G K E F L R E T W T R F M E D S F P I V N D Q E V M D V F L
SpliNPV2: V K P D T M K L I V N W N G K E F L R E T W T R F M E D S F P I V N D Q E V M D V F L
SpliNPV3: V K P D T M K L I V N W N G K E F L R E T W T R F M E D S F P I V N D Q E V M D V F L
SpliNPV4: V K P D T M K L I V N W N G K E F L R E T W T R F M E D S F P I V N D Q E V M D V F L

SpliNPVa: V V N M R P T R P N R C F R F L A Q H A L R C D P E Y V P H D V I R I V E P S Y V G T N
AcMNPV: V I N M R P T R P N R C F R F L A Q H A L R C D P D Y V P H E V I R I V E P V Y V G N N
SpliNPV1: V V N M R P T R P N R C F R F L A Q H A L R C D P E Y V P H D V I R I V E P S Y V G T N
SpliNPV2: V V N M R P T R P N R C F R F L A Q H A L R C D P E Y V P H D V I R I V E P S Y V G T N
SpliNPV3: V V N M R P T R P N R C F R F L A Q H A L R C D P E Y V P H D V I R I V E P S Y V G T N
SpliNPV4: V V N M R P T R P N R C F R F L A Q H A L R C D P E Y V P H D V I R I V E P S Y V G T N

SpliNPV: N E Y R I S L A K K G G G C P V M N L H A E Y T T S F E S F I D K V I W Y N F Y K P I V Y I G T
AcMNPV: N E Y R I S L A K K G G G C P V M N L H S E Y T H S F E E F I N R V I W E N F Y K P I V Y V G T
SpliNPV1: N E Y R I S L A K K G G G C P V M N L H A E Y T T S F E S F I D K V I W Y N F Y K P I V Y V G T
SpliNPV2: N E Y R I S L A K K G G G C P V M N L H A E Y T T S F E S F I D K V I W Y N F Y K P I V Y V G T
SpliNPV3: N E Y R I S L A K K G G G C P V M N L H A E Y T T S F E S F I D K V I W Y N F Y K P I V Y V G T
SpliNPV4: N E Y R I S L A K K G G G C P V M N L H A E Y T T S F E S F I D K V I W Y N F Y K P I V Y V G T
```

Figure 4. Comparison of amino acid sequence of the putative polypeptide of Polh-cr (Acc# AAR04375) to its corresponding sequences of AcMNPV (Acc# AAA46736) and other 4 SpliNPVs (Acc# AAW49208, BAA00824, AAT10182 and AAW49207, respectively). The different amino acids were bolded and shaded.

Discussion

To date, genetically engineered baculoviruses introduce a promising research line to overcome the slow action of baculoviruses as biocontrol agents. On the other hand, searching for new natural baculovirus isolates with better insecticidal characteristics is still a developing subject of work (more safe and has not the risks of releasing genetically engineered product in nature). Although AcMNPV is considered as the type species in the genus nucleopolyhedrovirus, 15 species and 471 tentative species of NPV have been isolated (26). Abul-Nasr (27) isolated an Egyptian NPV from the cotton leaf worm *S. littoralis*. Cherry and Summers (28) isolated the two reference NPV types A and B. During 1986- 1988, six natural isolates of SpliNPV were isolated from Giza, Menya, Kaha, Tokh, Kafr-Elsheikh and Gharbyia, Egypt (Khamiss, personal communication). Also, fifteen isolates of SpliNPV were isolated from the six above mentioned Egyptian localities in addition to Fayoum, Sakkara, Menofya, Benisuef, Asyut, Sinnuris, Banha, Sharkyia and Elsaff-Giza, Egypt (Khamiss, personal communication).

Combining our data with that presented by Khamiss and by Seufi (personal communication), it can be concluded that 34 NPV isolates have been collected from Egypt between 1986 and 2000. This may reflect the suitability of the Egyptian environment for not only gathering and collecting new baculovirus isolates but also for using them as potential pesticides in integrated pest management (IPM) programs. Given that baculoviruses have been isolated from Upper as well as Lower Egypt, in which the cultivation was washed by chemical insecticides, it would be expected to find more and more isolates in virgin regions (where no insecticides were used).

The susceptibility of 2nd instar larvae of *S. littoralis* to the two polyhedrovirus species reported in this study was comparable to that observed by Lacey et al. (21). They reported that LC₅₀s were 1.77×10^3 and 3.05×10^3 occlusion bodies (OB)/ mm² for *Anagrapha falcifera* (AfMNPV) and *Autographa californica* (AcMNPV) when applied to the neonate codling moth larvae. Our Egyptian isolate (SpliNPV) showed similar results to that obtained by Klein and Podoler against the Egyptian cotton leafworm (29). Abot et al. (30) clarified that LC₅₀s of an NPV isolate against two *A. gemmatalis* populations varied from 129 to 316 OBs/ ml diet. Their results fall within the range obtained with our isolate. On the other hand, our results showed lower LC₅₀ when compared with that reported by Pawar and Ramakrishnan (31) and Komolpith and

Ramakrishnan (32) whereas LC₅₀ was 4.677×10^6 PIB/ ml for 4-day old *S. littoralis* larvae. Similarly, Ashok and Ramakrishnan (33) reported higher LC₅₀ (7.1×10^6 PIB/ ml) for 3-day old *S. litura* larvae. Also, Stiles and Himmerich (34) introduced higher LC₅₀ of AcMNPV against *H. zea* ($3.46 \times 10^4 - 6.38 \times 10^5$ PIB/ ml). Finally, Abdel-Aziz (personal communication) presented higher LC₅₀s (1.8×10^7 and 9.0×10^7 PIB/ ml, respectively) for 2nd larval instar of *S. littoralis*. The variability of LC₅₀s is probably due to the method of surface treatment, homogeneously treated diet, feeding habit of the insect species (21) or due to difference in larval age (35,36). It may also be due to difference in host susceptibility to NPV (4), number of virions per occlusion body, virulence of the virus strain and/ or the difference in number of laboratory propagation cycles for the viral isolate. According to Van-Beek and Huges (37), the virulence of baculoviruses is best determined by the speed with which a given virus kills the insect pest.

LT values presented in Table I indicated that SpliNPV killed 2nd instar larvae of *S. littoralis* one day faster than AcMNPV. In comparing our results with that presented by other authors, many considerations have to be taken into account. Host range and LC₉₅ of the viral isolates are the most important considerations. Although LC₉₅ and LT₉₅ that produced in this study are economically prohibitive, improving the insecticidal characteristics of such isolates (by formulation and synergistic additives) is a growing subject in many companies.

The full length of polyhedrin gene from lepidopteran NPVs ranged from 483 bp to 747 bp (38). In case of polyhedrin gene from *Spodoptera* sp. NPVs, its full length ranges from 510 bp to 747 bp in comparison to that from *Autographa* sp. NPVs which ranges from 507 bp to 738 bp (38). Therefore, it could be said that Polh-cr represented about 65% of the full length of polyhedrin gene. On comparing nucleotide sequence of Polh-cr to all available sequences in the GenBank, it created a significant homology with 100 NPV and 11 GV genes. It showed 99% identity with *S. littoralis* polyhedrin gene (Acc# D01017), 95% with *S. littura* polyhedrin gene (Acc# AY552474) and 93% with *S. littura* polyhedrin genes (Acc# AY549963, AF325155, AF037262 and AF068189). In addition, it was 90% similar to *Lymantria dispar* polyhedrin genes (Acc# AF499687, AF081810 and M23176), 88% to *Malacosoma neustria* polyhedrin gene (Acc# X55658) and 87% to *B. mori* polyhedrin genes (Acc# M10043 and X63614). Furthermore, it showed

86% homology with *S. litura* and *Amsacta albistriga* polyhedrin genes (Acc# X94437 and AF118850, respectively) and 85% with *S. exigua* and *Malacosoma neustria* polyhedrin gene (Acc# AF169823; AY127899 and AJ277555, respectively). These results ensured that Polh-cr is a highly conserved region within polyhedrin gene of about one sixth of the known NPV species. Consequently, it could be used in many molecular techniques concerned with baculoviruses.

Knowing that the full length of polyhedrin protein from lepidopteran NPVs ranges from 161 a.a. (e.g. *Hyphantheria cunea* NPV, Acc# AAW49190 and *Bombyx mori*, Acc# ABB16300) to 249 a.a. (e.g. *Spodoptera litura* NPVs, Acc# AAZ78353, NP_258269 and AAS90121). In case of *Spodoptera* sp. NPV, full length of polyhedrin protein ranges from 170 a.a. (Acc# AAW49204) to 249 a.a. (Acc# AAZ78353) in comparison to *Autographa* sp. NPV which ranges from 169 a.a. (Acc# AAW63393) to 246 a.a. (Acc# AAA46736). The deduced amino acid sequence of Polh-cr was compared to other polyhedrins. Alignment results revealed that Polh-cr was 100% and 99% identical to *S. littoralis* polyhedrins (Acc# AAC33752, AAR04375; P24646, JU0382 and BAA00824, respectively). It was also 99% similar to *S. litura* polyhedrins (Acc# NP258269, AAC09246, AAL01689 and AAS58468). In addition, it has 98% identity with polyhedrin of *S. littoralis* (Acc# AA590121). Furthermore, it showed 89% similarity with polyhedrins of *S. exigua* and *Ecotropis obliqua* polyhedrins (Acc# AAF33532, JQ1868, NP037761, 001586; AAB53632, AAQ88174, P07388 and AAA46739).

Using AcMNPV nucleotide sequence (Acc# M25054) and amino acid sequence (Acc# AAA46736) as references for comparison with our sequences, it was found that 57

different nucleotides and 12 gaps in nucleotide sequence resulted in 15 different amino acids in the putative polypeptide (I, N, V, V, E, D, S, T, A, T, S, D, K, Y and I a.a. from our putative polypeptide were replaced with V, S, I, I, D, E, V, N, S, H, E, N, R, E and V a.a. from AcMNPV putative polypeptide, respectively). Surprisingly, Polh-cr putative polypeptide was identical to SpliNPV polypeptides (Acc# AAW49208, BAA00824, AAT10182 and AAW49207) with one amino acid replacement (I with V at the position 133). These results suggested the possible difference in codon usage among the compared isolates. It might also give a specific property to our putative polypeptide.

In this paper, we described the cloning and sequencing of a highly conserved region in polyhedrin gene and the insecticidal activity of an Egyptian NPV isolate. To our knowledge, this is the first report that determines the sequence of this conserved region in an Egyptian isolate. Further studies to develop kits for ELISA, Western and dot blotting, hybridization as well as potential biocontrol agent are switched on. The availability of Polh-cr products would be helpful in studies concerning ELISA, PCR and other related molecular techniques. In addition, it provides a candidate for effective, sensitive and reproducible diagnostic tools for screening insects or/ and other arthropods, especially crustacean species, crabs and shrimps for baculovirus infections and may be important in controlling (preventing/ enhancing) baculovirus infection. Also, it may be useful in monitoring the distribution of NPVs, the fate of genes and release of wild type as well as genetically engineered NPVs. Furthermore, it will facilitate risk assessment, ecological and viral epidemiological studies.

References

1. Martignoni ME, Iwai PJ. *A catalog of viral diseases of Insects, Mites and Ticks*, 4th edn. Portland, Oregon: USDA Forest Service PNW 1986; 195.
2. Huber J. Use of baculoviruses in pest management programs. In: "The biology of baculoviruses", (R. R. Granados and B. Federici, Editors), CRC Press, Boca Raton, FL. 1986; 2: 181-202.
3. Wood HA, Granados RR. Genetically engineered baculoviruses as agent for pest control. *Annu Rev Microbiol.* 1991; 45:69-87
4. El-Salamouny S, Lange M, Jutzi M, Huber J, Jehle JA. Comparative study on the susceptibility of cutworms (Lepidoptera: Noctuidae) to *Agrotis segetum* nucleopolyhedrovirus and *Agrotis ipsilon* nucleopolyhedrovirus. *J Invertebr Pathol.* 2003; 84: 75-82.
5. Wood HA. (1996): Genetically enhanced baculovirus insecticides. In "Molecular biology of the biological control of pest and diseases of plants" (M. Gunasekaran, DJ Weber Eds.), CRC Press, Boca Raton, 91-104.
6. Couch JA. Free and occluded virus, similar to baculovirus, in hepatopancreas of pink Shrimp. *Nature (London)* 1974;247: 229-231.
7. Hsu YL, Wang KH, Yang YH, Tung MC, Hu CH, Lo CF, Wang CH, Hsu T. Diagnosis of *penaeus monodon*-type baculovirus by PCR and by ELISA of occlusion bodies. *Dis Aquat Organ.* 2000; 40: 93-99.
8. Halder M, Ahne W, Thomsen I. Detection of a baculovirus in the tiger prawn *Penaeus monodon*. *Zentralbl Veterinarmed B.* 1989; 36: 257-260.
9. Chang PS, Lo CF, Kou GH, Lu CC, Chen SN. Purification and amplification of DNA from *Penaeus monodon*-type baculovirus (MBV). *J Invertebr Pathol.* 1993; 62: 116-120.
10. Poulos BT, Mary J, Bonami JR, Redman R, Lightner DV. Use of non-radioactively labeled DNA probes for the detection of a baculovirus from *Penaeus monodon* by in situ hybridization of fixed tissue. *J Virol Methods.* 1994; 49: 187-193.
11. Faktor O, Raviv D. A polymerase chain reaction for the detection of nucleopolyhedroviruses in infected insects: the fate of the *Spodoptera littoralis* virus in *Locusta migratoria*. *J Virol Methods.* 1996; 61: 95-101.
12. Belcher CR, Young PR. Colourimetric PCR-based detection of monodon baculovirus in whole *Penaeus monodon* postlarvae. *J Virol Methods.* 1998; 74: 21-29.
13. Wang CH, Yang HN, Liu HC, Kou GH, Lo CF. Nested polymerase chain reaction and in situ hybridization for detection of nucleopolyhedrosis. *J Virol Methods.* 2000; 84: 65-75.
14. Christian PD, Gibb N, Kasprzak AB, Richards A. A rapid method for the identification and differentiation of *Helicoverpa nucleopolyhedroviruses* (NPV Baculoviridae) isolated from the environment. *J Virol Methods.* 2001; 96: 51-65.
15. De Moraes RR, Maruniak JE. Detection and identification of multiple baculoviruses using the polymerase chain reaction (PCR) and restriction endonuclease analysis. *Mol Cells.* 2001; 11: 334-340.
16. Ebling PM, Smith PA, van Frankenhuyzen K. DNA hybridization assay for detection of nucleopolyhedrovirus in whitemarked tussock moth (*Orgyia leucostigma*) larvae. *Pest Manag Sci.* 2001; 57: 66-71.
17. Woo SD. Rapid detection of multiple nucleopolyhedroviruses using polymerase chain reaction. *Mol Cells.* 2001; 11: 334-40.
18. Murillo R, Munoz D, Williams T, Mugeta N, Caballero P. Application of the PCR-RFLP method for the rapid differentiation of *Spodoptera exigua* nucleopolyhedrovirus genotypes. *J Virol Methods.* 2006; 135: 1-8.
19. El-Defrawi ME, Tappozada A, Mansour N, Zeid M. Toxicological studies on Egyptian cotton leaf worm *Prodenia litura* (F.)1: susceptibility of different larval instars of *Prodenia litura* to insecticides. *J Econ Entomol.* 1964; 57: 591-593.
20. Levinson ZH, Navon A. Ascorbic acid and unsaturated fatty acids in the nutrition of the Egyptian cotton leaf worm. *J Insect Physiol.* 1969; 15: 591-595.

21. Lacey LA, Vail PV, Hoffmann DF. Comparative activity of baculoviruses against the codling moth *Cydia pomonella* and three other tortricid pests of tree fruit. *J Invertebr Pathol.* 2002; 80: 64-68.
22. Abbott WS. A method of computing the effectiveness of an insecticide. *J Econ Entomol.* 1925; 18: 265-267.
23. Throne JE, Weaver DK, Chew V, Baker JE. Probit analysis of correlated data: multiple observations over time at one concentration. *J Econ Entomol.* 1995; 88: 1510-1512.
24. Saiki RK, Gelfand DH, Stoffel S, Scharf SJ, Higuchi R, Horn GT, Mullis KB, Erlich HA. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 1988; 239: 487-491.
25. Abdel-Hamid I, Salama MS, Madkour MA. The use of universal nested polymerase chain reaction (UN-PCR) to detect insect nuclear polyhedrosis viruses. *J Egypt Ger Soc Zool.* 1996;20(E): 97-107.
26. Volkman LE, Blissard GW, Friesen P, Keddie BA, Possee R, Theilmann DA. Family baculoviridae. In : *Virus taxonomy: Classification and nomenclature of viruses. Sixth report of the International committee on taxonomy of viruses.* Murphy FA, Fauquet CM, Bishop DHL, Ghabrial SA, Jarvis AW, Martelli GP, Mayo MA, Summers MD, Editors. Vienna: Springer- Verlag; 1995. p104-113.
27. Abul-Nasr S. Further tests on the use of a polyhedrosis virus in the control of the cotton leaf worm *Prodenia litura*. *J Insect Pathol.* 1959; 1: 112-120.
28. Cherry CL, Summers MD. Genotypic variation among wild isolates of two nuclear polyhedrosis viruses isolated from *Spodoptera littoralis*. *J Invertebr Pathol.* 1985; 46: 289-295.
29. Klein M, Podoler H. Studies on the application of a nuclear polyhedrosis virus to control population of the Egyptian Cotton worm, *Spodoptera littoralis*. *J Invertebr Pathol.* 1978; 32: 244-248.
30. Abot AR, Moscardi F, Fuxa JR, Sosa-Gomez DR, Richter AR. Susceptibility of populations of *Anticarsia gemmatalis* (Lepidoptera: Noctuidae) from Brazil and the United States to a nuclear polyhedrosis virus. *J Entomol Sci.* 1995;30: 62-69.
31. Pawar VM, Ramakrishnan N. Investigation of the nuclear polyhedrosis virus of *Spodoptera litura* (Fab). III. Bioassay of virus activity. *Indian J Entomol.* 1975; 37: 374-376.
32. Komolpith U, Ramakrishnan N. Bioassay of the nuclear polyhedrosis virus against larval stages of *Spodoptera litura* (Fab) and the effect of protectants against ultraviolet light. *Proc Indian Acad Sci.* 1975; 83: 195-203.
33. Ashok MV, Ramakrishnan N. Certain observations on the physico-chemical properties of nuclear polyhedrosis virus of *Spodoptera littoralis* (Fabricius). *J Entomol Res.* 1981; 5: 43-46.
34. Stiles B, Himmerich S. *Autographa californica* NPV isolates: Restriction endonuclease analysis and comparative biological activity. *J Invertebr Pathol.* 1998; 72: 174-177.
35. Pyne CC. Insect viruses as control agents. *Parasitol.* 1982; 84: 35-77.
36. Pyne CC, Tatchell GM, Williams CF. The comparative susceptibilities of *Pieris brassicae* and *Pieris rapae* to a granulosis virus from *Pieris brassicae*. *J Invertebr Pathol.* 1981; 38: 273-280.
37. Van Beek NA, Hughes PR. The response time of insect larvae infected with recombinant baculoviruses. *J Invertebr Pathol.* 1998; 72: 338-347.
38. Jehle JA, Lange M, Wang H, Hu Z, Wang Y, Hauschild R. Molecular identification and phylogenetic analysis of baculoviruses from Lepidoptera. *Virol.* 2006; 346: 180-193.